



Studies on the antimicrobial potency of the marine algae - *Caulerpa scalpelliformis* and *Spyridia* species

Mrinalini.J.Singh

Research Scholar, Department of Botany, Nirmala College for Women, Coimbatore 641 018, Tamil Nadu, India

E-mail: mrinalini.singh50@gmail.com

Abstract

Objective: To study the antimicrobial potency of the marine algae – *Caulerpa scalpelliformis* and *Spyridia* sp.

Methods: *Caulerpa scalpelliformis* and *Spyridia* sp. were screened for the presence of bioactive components by extracting them with suitable solvents and studying their antimicrobial activities against varied human pathogens. An attempt has also been made to separate the unknown compounds using Thin Layer Chromatography and purify the extracts using column chromatography since not much work has been carried out in these species.

Results: In *Spyridia*, only the methanolic extract could produce zone of inhibition against *Staphylococcus aureus* and *Salmonella* sp. whereas in *Caulerpa scalpelliformis*, only the petroleum ether, chloroform and methanolic extracts showed antimicrobial properties. Better resolution was found in the mobile phase containing chloroform: methanol 80: 20 ratio for *Caulerpa*. Hence, its methanolic extract was fractionated in silica gel column and these fractions were effective against some of the test organ-isms but not against all.

Conclusion: Further investigation is required to study the nature of compounds with antimicrobial properties with reference to *Caulerpa scalpelliformis*; whereas with *Spyridia* sp. the study can be extracted to collect the samples during different seasons of a year and confirm its antimicrobial potency.

Keywords: *Caulerpa Scalpelliformis*, *Spyridia* SP, Antimicrobial Activities, Thin Layer Chromatography, Column Chromatography.

1. Introduction

Marine algae or seaweeds are salt water dwelling, containing various pigments and are attached by hold fasts to anchor them. They are rich in bioactive natural products and have been studied as potential biocides and pharmaceutical agents (Abbott et al., 2002). Among the seaweeds group, the members of green and red marine algae are of great economic importance. The main products of red seaweeds are agar-agar and carrageen used mostly in baking, and as an emulsifying agent in pharmaceutical industry.

Due to the emergence of commercial drug resistant organisms with enhanced side effects, there is a need for the discovery of new drugs. Marine algae seem to be the treasure house of natural bioactive secondary metabolites with high therapeutic potency and no side effects. Despite this, the number of seaweed species studied and identified corresponds to only 2% of the 150,000 known species worldwide (Harvey et al., 1988). From the species identified, approximately two thousand chemical compounds have been characterized (González & Silva, 2001). Various systematic works have been carried out on marine algal extracts worldwide for antimicrobial, anticancerous, anti-inflammatory and antioxidant studies with no side effects.

Hence, the objective of this study was to evaluate the antibacterial activities of the organic extracts from two marine algae *Caulerpa* sp and *Spyrida* sp collected from the South Eastern coastal areas of Tamil Nadu.

2. Materials and methods

2.1. Sample collection

About 1 kg of the two species of seaweeds namely *Caulerpa* sp and *Spyrida* sp were collected fresh from the south east coast of Tamil Nadu (Rameshwaram) during the month of December, 2007. Cleaning and drying of the samples were performed as mentioned by Pereira et al., 2003. The washed samples were shade dried, powdered and stored in sterile containers under refrigeration until use.

Extraction of hundred grams of powdered biomass with solvents of increasing polarity ranging from petroleum ether to water through benzene, chloroform, ethyl acetate and methanol for 72 hours at room temperature with intermittent stirring for every twenty four hours was made in succession.

Strains used for testing antimicrobial activity were *Staphylococcus aureus* (MTCC, 740), *Salmonella typhimurium* (MTCC, 98), *Escherichia coli* (GM242) and *Candida albicans* (MTCC, 227). The remaining isolates such as *Klebsiella* sp, *Proteus* sp, *Citrobacter* sp and *Pseudomonas* sp were collected from PSG Institute of Medical Science & Research, Coimbatore.

Around 100 µl of all the extracts thus were injected into empty sterilized filter paper disc having a diameter of 5 mm. The discs impregnated with the mother solvents of each extracts served as the control and were placed on the same plate. Antimicrobial activities of algal extracts were tested separately using disc diffusion method (Bauer et al., 1966).

Thin layer chromatography of each sample was performed on Merck TLC F254 plates, with Chloroform: Methanol in the ratio

of varying concentrations such as 10: 90, 80:20, 20:80 and 90:10 as mobile phase. The separated components were visualized under ultraviolet light of 254 nm.

Purification of the crude extracts was carried out as per the procedure of Vairappan *et al.*, 2001 with few modifications. The seaweed extracts with antibacterial activity were fractionated by silica gel column chromatography (chloroform and methanol). The fractions were eluted with chloroform: methanol and were further subjected to antimicrobial assay and Thin layer chromatography.

3. Result

Table 1: Antimicrobial Activity from Various Extracts of *Spyridia* Sp.

Test Organisms	Zone of Inhibition (mm)				
	Petroleum ether	Benzene	Chloroform	Ethyl acetate	Methanol
<i>Escherichia coli</i>	-	-	-	-	-
<i>Staphylococcus aureus</i>	-	-	-	-	7.5
<i>Citrobacter</i> sp.	-	-	-	-	-
<i>Klebsiella</i> sp.	-	-	-	-	-
<i>Pseudomonas</i> sp.	-	-	-	-	-
<i>Proteus</i> sp.	-	-	-	-	-
<i>Salmonella</i> sp.	-	-	-	-	9.4
<i>Candida albicans</i>	-	-	-	-	-

Key: '-' sign indicates the absence of zone of inhibition

Table 2: Antimicrobial Activity from Various Extracts of *Caulerpa scalpelliformis*

Test Organisms	Zone of Inhibition (mm)				
	Petroleum ether	Benzene	Chloroform	Ethyl acetate	Methanol
<i>Escherichia coli</i>	8	-	10	-	8
<i>Staphylococcus aureus</i>	-	-	-	-	-
<i>Citrobacter</i> sp.	-	-	8	-	10
<i>Klebsiella</i> sp.	9	-	-	7	8
<i>Pseudomonas</i> sp.	-	-	-	8	8
<i>Proteus</i> sp.	-	-	-	-	-
<i>Salmonella</i> sp.	-	-	-	-	-
<i>Candida albicans</i>	-	-	-	-	9

In order to survive in a highly competitive environment marine algae have to develop defense strategies that results in a tremendous diversity of compounds from a different metabolic pathway. Recent trends in drug research from natural sources have shown that the algae are promising organisms to furnish novel biochemically active compounds. The bioactive substances from seaweed have great chances to be employed against bacteria, viruses, other pathogens and tumors (Cragg *et al.*, 1997).

The crude extracts were used to study their antimicrobial properties with the test organisms. In *Spyridia*, only the methanolic extract could produce zone of inhibition against *Staphylococcus aureus* and *Salmonella* sp. (Table 1) whereas in *Caulerpa scalpelliformis*, only the petroleum ether, chloroform and methanolic extracts of *Caulerpa scalpelliformis* showed antimicrobial properties (Table 2).

Thin layer chromatography was performed to find out a better separation of compounds using various combinations of chloroform and methanol and a better resolution was found in the mobile phase containing chloroform: methanol 80: 20 ratio. The methanolic extract of *Caulerpa scalpelliformis* was fractionated in silica gel column. The elutions were collected at regular intervals

and were checked for their antimicrobial properties. The fractions were effective against some of the test organisms but not against all (Table 3).

Table 3: Antimicrobial Activity of Partially Purified Methanol Extract through Silica Gel Column Chromatography from *Caulerpa Scalpelliformis*

Test organisms	Zone of inhibition(mm)								
	1	2	3	4	5	6	7	8	9
<i>E. coli</i>	-	10	11	10	10	-	-	10	-
<i>Staph. aureus</i>	10	10	-	-	-	10	11	10	10
<i>Citrobacter</i> sp.	9	8	8	-	8	10	-	-	-
<i>Klebsiella</i> sp.	-	-	-	10	8	8	8	-	8
<i>Pseudomonas</i> sp.	-	-	-	-	-	-	-	10	8
<i>Proteus</i> sp.	-	-	-	9	-	-	-	-	-
<i>Salmonella</i> sp.	-	-	-	-	-	-	-	9	-
<i>Candida albicans</i>	10	9	-	-	10	-	-	-	-

In the crude methanolic extract, effect was observed against all the test cultures whereas none of the eluted fractions could inhibit few test cultures.

4. Discussion

The probable reasons attributed to this are one single compound was not sufficient to inhibit the growth of these cultures. This could further be supported but the presence of antimicrobial compounds in various fractions. The genus *Caulerpa* is represented by nearly 60 species. Several reports are available on the metabolic of various other species of *Caulerpa*. No reports in this regard are available on the antibacterial compounds of *C. scalpelliformis*. Hence further purification and structure analysis only could answer the above mystery of none of the fractions inhibiting few test cultures.

Finally it could be concluded that the *C. scalpelliformis* is a potential source of natural antibiotics.

5. Conclusion

In this paper, we have reported the antimicrobial properties of bioactive components extracted from the marine algae *Caulerpa scalpelliformis* and *Spyridia* sp. Further investigation is required to study the nature of compounds with antimicrobial properties with reference to *Caulerpa scalpelliformis*; whereas with *Spyridia* sp. the study can be extracted to collect the samples during different seasons of a year and confirm its antimicrobial potency. Intensive studies must be carried out to unravel its unexhausted reserve of bio-metabolites.

Conflict of Interest Statement

We hereby declare that we have no conflict of interest.

Acknowledgement

I am thankful to the authorities of Microbiology Department especially Late Mrs. C.K. Raadha in PSG College of Arts and Science, Coimbatore, Tamil Nadu for providing necessary facilities to perform the experiment.

References

- [1] Abbott, I.A., Fisher, J. Mc Dermid, K.J. (2002). New re-reported and revised marine algae from the Nha Trang, Vietnam, in: taxonomy of economic seaweed with reference to some pacific species. Oceanographic Institute Nha Trang Vietnam 8:291-321
- [2] Harvey SC, J Luo & R Lavery. (1988). DNA stem-loop structures in oligopurine-oligopyrimidine triplexes. Nucleic Acids Research 16(24): 11795-11809. <http://dx.doi.org/10.1093/nar/16.24.11795>.
- [3] González F & M Silva. (2001). Biodiversidad química de macroalgas marinas. In: Alveal K & T Antezana (eds). Sustentabilidad de la biodiversidad, un problema actual. Bases científico-técnicas, teorizaciones y proyecciones, and pp. 415-496. Universidad de Concepción, Concepción.
- [4] Pereira R.C., B.A.P. DaGama, V.L. Teixeira and Y. Yoneshigue-Valentín (2003). Ecological roles of natural products of the Brazilian red seaweed *Laurencia obtuse*. Braz.J.Biol 63(4):665-672. <http://dx.doi.org/10.1590/S1519-69842003000400013>.
- [5] Bauer, A.W., W.M. Kirby, J.C. Scheris and M. Truck (1966). Antibiotic susceptibility testing by standardization by single disc method. Amer.J. Clinical Pathol 45:493-496.
- [6] Vairappan C. S., Suzuki, M., Abe T., Masuda, M. (2001). Halogenated metabolites with antibacterial activity from the Okinawa, *Laurencia* sp phytochemistry 58: 517-523.
- [7] Cragg, G. M., Newman and K. M. Snader, (1997). Natural products in drug discovery and development J.Nat prod 60: 52-60. <http://dx.doi.org/10.1021/np9604893>.