

Impact on post partum amenorrhoea beyond breastfeeding

A. K. Shukla ^{1*}, A. K. Tiwari ², R. C. Yadava ³

¹ Assistant Professor Department of Community Medicine Subharti Medical College
Swami Vivekanad Subharti University, Meerut-250005 (UP) India

² Faculty Member IBS Business School ICFAI University Dehardun India

³ Department of Statistics Banaras Hindu University (BHU) Varanasi India

*Corresponding author E-mail: arvindshukla_vns@rediffmail.com

Abstract

Citing various studies Bongaarts (1980) has concluded that breastfeeding is the principal determinant of post partum amenorrhoea (PPA) and unrestricted breastfeeding is associated with lower fertility. On the other hand many researchers have shown opposite views by saying that other factors are also responsible in affecting the duration of PPA. In this paper, we have investigated the relationship between the two duration variables viz. PPA and breastfeeding utilizing data of NFHS-2. For this purpose, we have controlled the duration of breastfeeding and then investigated whether distribution of PPA duration shows similar pattern in different situations or it varies with the situations. This can help in deciding whether PPA duration is mainly affected by breastfeeding or some other factors are also operating in shaping its distribution. In this context, we show that even after controlling the duration of breastfeeding (keeping breastfeeding at least six months) the percentage of females having PPA duration less than four months, shows large variation according to caste, educational level, standard of living index (SLI) and body mass index (BMI) of females. If breastfeeding would have been the only principal determinant of duration of PPA, this percentage should have remained almost constant! This shows that some other factors beyond breastfeeding are responsible for shaping the distribution of PPA duration of a female.

Keywords: Post Partum Amenorrhoea (PPA); Body Mass Index; NFHS; SLI.

1. Introduction

The duration of post partum amenorrhoea (PPA) is the period taken to return of menses following a live birth or late term of abortion or still birth? This is temporary infecundable period during which a woman is non-susceptible to pregnancy again due to suspension of natural cycle of ovulation and menstruation and the attainment of first menstruation is treated as the termination of PPA.

Fertility is basically a biological phenomenon very much influenced by a large number of socio-cultural and behavioral factors. This fact greatly complicates the analysis of the effect of the biological factors on fertility.

Among the many biological factors affecting human fertility, duration of PPA plays a dominant role in shaping the fertility behavior of a population. It is extensively reported that breastfeeding is the major determinant of prolonged PPA, in societies where it is nearly universal, prolonged, and of high intensity (Santow, 1987; Howie and McNeilly, 1982; Habicht et al., 1985; Akin et al., 1986; Huffman et al., 1987; Srinivasan et al., 1989; Savina and Kennedy, 1989; Nath et al., 1993; Singh et al., 1994). Many researchers have pointed out that breastfeeding is the only dominant factor influencing the PPA and especially nutrition does not play any significant role on it (Bongaarts, 1980). This conclusion is mainly based on the results obtained in Choudhary (1978), Huffman et al. (1978) Bongaarts and Delgado (1979) etc. It is because of this fact that Bongaarts (1978) has considered duration of breastfeeding as one of the important proximate determinants of fertility.

On the other hand, opposite views have been reported by many other researchers (Chebez and Maliner, 1973; Huffman et al.,

1987; Prema et al., 1981; Lune et al., 1981; ICMR 1995; Diezet et al., 1998; Yadav et al., 2005). The authors have shown that better nutrition is responsible for shortening the duration of PPA. Many researchers have shown that several socio-economic, demographic and cultural factors affect the length of PPA (Delgado et al., 1982; Habicht et al., 1985; Singh and Singh, 1989; Singh et al., 1990; Nath et al., 1993; Singh et al., 1994). Most of these results are based on data of limited areas and sample sizes are also not reasonably large.

In the National Family Health Survey (NFHS), extensive data have been collected on durations of post partum amenorrhoea (PPA) as well as breastfeeding for births occurring during last three or four years. Such data provide great opportunities to investigate the relationship between the two duration variables. Most of the studies have shown that there is strong positive relationship between two variables. However, still the view is quite open whether duration of breastfeeding is the only dominant variable influencing PPA or there are some other factors also which play significant role on determining PPA. The objective of the present paper is to throw some light on this complicated issue. For this purpose data of NFHS-2 for various major states have been utilized to investigate the nature of relationship as well as variation (if any) in the spatial pattern in the country.

2. Data

The National Family Health Survey (NFHS) was carried out as the principal activity of a collaborative project to strengthen the research capabilities of the Population Research Centers in India, initiated by the Ministry of Health and Family Welfare, Govern-

ment of India, and coordinated by the International Institute for Population Sciences, Mumbai. There have been two rounds of the survey conducted. (Recently the third round of NFHS has also been completed but detailed results are still awaited). The first round of NFHS was conducted in 1992-93 (NFHS-1) while second round was undertaken in 1998-99 (NFHS-2). In the NFHS-2, data relating to current status of post partum amenorrhoea related to births occurring during last three years from the survey date have been obtained. Further, for the cases where post partum amenorrhoea is already completed, the duration of PPA have also been noted. These data constitute the basis of study in this paper.

3. Methodology and results and discussion

The NFHS report has already provided estimates of median duration of breastfeeding as well as PPA period for different states of the country. These are presented here as table 1.

A look at this table clearly indicates that there is strong positive relationship between average (median) durations of breastfeeding and PPA. Is this much analysis adequate to conclude that duration of breastfeeding is a dominant factor affecting duration of PPA? The answer is perhaps "not".

We try to investigate the matter here in some alternative way. Such attempt has also been done by Yadav et. al. (2005). But the results are based on data from a very limited area and the sample size is also small. The whole idea behind this type of analysis is that if we fix the duration of breastfeeding and examine the behavior of duration of PPA, the results may give some clues. Consequently we have fixed the duration of breastfeeding (at least six months) and then compare the proportion of births where duration of PPA is less than four months. For all such cases, where the PPA period is ceased, breastfeeding was still continuing and all births have occurred at least six months prior to survey date. Thus if breastfeeding is the only dominant factor affecting PPA period, then these proportions should be almost constant for all groups of the population.

We have computed such proportions (in percentages) for different groups of the major states of the country classified according to caste category, religion, area of residence, educational level, body mass index (BMI) and standard of living index (SLI).

These results are shown in table 2 to 15 for states of Uttar Pradesh, Madhya Pradesh, Rajasthan, Bihar, Orissa, Punjab, Tamil Nadu, Kerala, Karnataka, Andhra Pradesh, Maharashtra, Gujarat, West Bengal and India as a 'whole'.

A close look on these tables reveals interesting and curious results. First of all there is large variation in this proportion across different groups in all the states of the country, the lowest being less than 15% (Karnataka and Andhra Pradesh) to maximum being near 70% (Punjab). We must remember that these estimated percentages are not based on smaller samples.

The other important feature can be seen from the following fact: Definitely standard of living index (SLI), body mass index (BMI) and educational level of females can be considered to be high correlates of social, economic and nutritional development of females under consideration. When we observe the computed percentages for different groups it is found that

- i) As BMI increases, by and large the proportion also increases constantly in all states.
- ii) As SLI increases the above pattern emerges for all the states.
- iii) Similar results are also found when we consider the educational level of females.
- iv) Similar situation also occurs when we observe the percentages according to caste groups. It is minimum for SC/ST in all the states where as the maximum is observed for 'Other caste'. The OBC's lie in between the two. Similar is the position of these groups as for as their social, economic and nutritional status are concerned.

- v) When we observe the rural-urban differentials in this percentage, the lower values are observed for rural areas in all the states.

These results clearly indicate that better economic and nutritional development is responsible for shortening the duration of PPA because increase in the percentage of PPA period less than four months will be responsible for lowering the average period of PPA.

These results have been obtained on the condition that everywhere duration of breastfeeding is more than six months. This implies that breastfeeding has no role in altering these proportions. Thus the obvious conclusion may be that some factor (s) beyond breastfeeding is/are playing significant role on shaping the behavior of PPA period. From the above discussion it seems that perhaps nutrition may be a dominant determinant of this proportion because BMI is considered to be a good indicator of nutritional status of a female.

The other important finding is that although the above pattern is almost universal but the economically most developed state of Punjab shows very high proportions (percentages) in comparison to less developed states like Orissa, Uttar Pradesh and Bihar for the same groups. For example, in Punjab for BMI>25.0, this percentage is 71.4 where as in the same group for Orissa, Uttar Pradesh and Bihar this is around 50%. The reasons are not very much clear but perhaps better nutritional status in this group in Punjab in comparison to other states may be responsible for such a pattern or some other factors such as food habits or other environmental or behavioral factors may be responsible for this phenomenon. This needs further investigations.

We have also analyzed the data according to religion. Only the two religions have been considered as Hindu and Muslim (except in Punjab where the two religions are Hindu and Sikh). In this case no uniform pattern is seen over the states. However, in majority of states, Hindus show lower percentage for PPA period less than four months in comparison to Muslims (India, Kerala, Tamil Nadu, Punjab, Gujarat, Maharashtra, Karnataka, Andhra Pradesh, Madhya Pradesh and Bihar). The opposite trend is observed for Rajasthan and Orissa. However the percentage for Hindus and Muslims are almost equal in case of Uttar Pradesh and West Bengal. In Punjab, Hindus show higher percentage than Sikhs.

4. Conclusion

Bongaarts (1980) concluded that breastfeeding was the principal determinant of PPA, and unrestricted breastfeeding was associated with lower fertility. The results obtained in this paper clearly indicate that even after controlling the duration of breastfeeding (keeping breastfeeding at least six months) the percentage of females having PPA duration less than four months, shows large variation according to caste educational level, SLI and BMI of female. This clearly indicates that some other factors beyond breastfeeding are responsible for shaping the distribution of PPA duration of a female.

Table 1: Median Duration of Breastfeeding and Post Partum Amenorrhoea (PPA) by States, NFHS-2, (1998-1999)

States	Median duration of Breastfeeding	Median duration of PPA
Kerala	24.5	4.0
Tamil Nadu	16.1	5.1
Punjab	21.2	5.4
West Bengal	≥36	6.7
Gujrat	22.0	7.5
Maharashtra	23.8	7.7
Karnataka	20.0	8.0
Rajasthan	25.5	8.5
Orissa	≥36	8.7
Uttar Pradesh	25.8	9.0
Andhra Pradesh	25.0	9.5
Madhya Pradesh	≥36	10.3
Bihar	≥36	11.3
INDIA	25.4	8.6

Table 2: Percentage of Females Whose PPA Less Than Four Months (and Breastfeeding Duration Is More Than Six Months) and Median Duration of PPA According to Different Characteristics, **INDIA**, NFHS-2

Characteristics	Percentage of females (whose PPA<4months, when BF>6)	Median duration of PPA (in months)
Total	29.2	8.2
Caste		
SC/ST	23.3	10.2
OBC	29.0	8.2
Others	34.5	6.6
Religion		
Hindu	28.2	8.3
Muslim	32.2	6.9
Place of Residence		
Rural	26.6	8.7
Urban	38.3	6.0
Education		
Illiterate	24.1	10.8
Literate (less than middle)	30.5	6.8
Middle to high school	36.2	6.2
Above high school	44.5	4.4
Body Mass Index		
<18.5	24.9	10.1
18.5-20.0	27.9	8.3
20.0-25.0	34.6	6.5
>25.0	46.9	4.3
Standard of Living Index		
Low	21.3	12.1
Medium	29.9	6.9
High	45.8	4.4

Table 3: Percentage of Females Whose PPA Less Than Four Months (And Breastfeeding Duration Is More Than Six Months) and Median Duration of PPA According To Different Characteristics, **Kerala**, NFHS-2

Characteristics	Percentage of females (whose PPA<4months, when BF>6)	Median duration of PPA (in months)
Total	45.9	4.5
Caste		
SC/ST	38.1	6.3
OBC	53.5	2.9
Others	49.4	4.1
Religion		
Hindu	45.9	4.5
Muslim	50.5	3.0
Place of Residence		
Rural	49.4	4.0
Urban	51.6	3.0
Education		
Illiterate	32.0	8.3
Literate (less than middle)	45.7	4.6
Middle to high school	54.9	2.9
above high school	52.2	2.9
Body Mass Index		
<18.5	43.8	3.9
18.5-20.0	49.3	4.1
20.0-25.0	54.5	2.9
>25.0	51.8	3.0
Standard of Living Index		
Low	41.1	6.1
Medium	25.3	6.9
High	58.8	2.8

Table 4: Percentage of Females Whose PPA Less Than Four Months (And Breastfeeding Duration Is More Than Six Months) and Median Duration of PPA According to Different Characteristics, **Tamil Nadu**, NFHS-2

Characteristics	Percentage of females (whose PPA<4months, when BF>6)	Median duration of PPA (in months)
Total	33.8	6.2
Caste		
SC/ST	26.7	6.7
OBC	36.1	6.0
Others	54.4	2.8
Religion		
Hindu	32.7	6.2
Muslim	42.8	4.5
Place of Residence		
Rural	29.9	6.4
Urban	41.7	4.7
Education		
Illiterate	28.0	6.7
Literate (less than middle)	32.2	6.5
Middle to high school	38.6	4.6
Above high school	43.9	4.4
Body Mass Index		
<18.5	30.7	6.4
18.5-20.0	25.9	6.3
20.0-25.0	37.0	6.1
>25.0	51.1	3.0
Standard of Living Index		
Low	25.5	6.8
Medium	37.6	6.1
High	47.7	4.1

Table 5: Percentage of Females Whose PPA Less Than Four Months (And Breastfeeding Duration Is More Than Six Months) and Median Duration of PPA According to Different Characteristics, **Punjab**, NFHS-2

Characteristics	Percentage of females (whose PPA<4months, when BF>6)	Median duration of PPA (in months)
Total	59.4	2.7
Caste		
SC/ST	52.0	2.9
OBC	58.6	2.7
Others	65.1	2.6
Religion		
Hindu	57.9	2.8
Sikh	51.8	2.9
Place of Residence		
Rural	57.8	2.7
Urban	64.2	2.4
Education		
Illiterate	54.6	2.8
Literate (less than middle)	50.4	3.0
Middle to high school	57.4	2.7
Above high school	69.4	2.3
Body Mass Index		
<18.5	47.6	4.3
18.5-20.0	61.3	2.7
20.0-25.0	61.2	2.6
>25.0	71.4	2.4
Standard of Living Index		
Low	42.4	4.6
Medium	51.8	2.9
High	67.8	2.4

Table 6: Percentage of Females Whose PPA Less Than Four Months (And Breastfeeding Duration Is More Than Six Months) and Median Duration of PPA According to Different Characteristics, **West Bengal**, NFHS-2

Characteristics	Percentage of females (whose PPA<4months, when BF>6)	Median duration of PPA (in months)
Total	28.5	8.0
Caste		
SC/ST	23.4	8.7
OBC	26.4	8.8
Others	31.8	6.8
Religion		
Hindu	28.8	6.8
Muslim	26.3	8.6
Place of Residence		
Rural	25.0	8.6
Urban	45.4	4.3
Education		
Illiterate	22.0	12.0
Literate (less than middle)	32.6	6.7
Middle to high school	33.3	6.4
Above high school	46.9	4.1
Body Mass Index		
<18.5	23.4	8.9
18.5-20.0	31.7	8.1
20.0-25.0	35.5	6.0
>25.0	41.7	4.5
Standard of Living Index		
Low	21.6	12.0
Medium	33.1	6.4
High	51.3	3.0

Table 7: Percentage of Females Whose PPA Less Than Four Months (And Breastfeeding Duration Is More Than Six Months) and Median Duration of PPA According to Different Characteristics, **Gujarat**, NFHS-2

Characteristics	Percentage of females (whose PPA<4months, when BF>6)	Median duration of PPA (in months)
Total	32.2	6.8
Caste		
SC/ST	26.5	10.6
OBC	28.7	8.1
Others	39.7	4.8
Religion		
Hindu	31.0	7.0
Muslim	43.0	4.4
Place of Residence		
Rural	26.5	9.0
Urban	41.5	4.8
Education		
Illiterate	22.8	12.0
Literate (less than middle)	34.9	8.3
Middle to high school	38.3	6.0
Above high school	49.4	4.0
Body Mass Index		
<18.5	24.5	10.6
18.5-20.0	30.5	8.7
20.0-25.0	42.7	4.8
>25.0	39.4	6.2
Standard of Living Index		
Low	19.7	12.4
Medium	33.1	6.7
High	44.6	4.4

Table 8: Percentage of Females Whose PPA Less Than Four Months (And Breastfeeding Duration Is More Than Six Months) and Median Duration of PPA According To Different Characteristics, **Maharastra**, NFHS-2

Characteristics	Percentage of females (whose PPA<4months, when BF>6)	Median duration of PPA (in months)
Total	24.5	8.6
Caste		
SC/ST	20.3	8.5
OBC	21.9	8.3
Others	28.0	7.0
Religion		
Hindu	24.3	8.4
Muslim	28.6	6.6
Place of Residence		
Rural	19.6	9.0
Urban	22.5	6.4
Education		
Illiterate	19.6	12.1
Literate (less than middle)	23.9	8.4
Middle to high school	23.8	7.0
above high school	36.8	6.0
Body Mass Index		
<18.5	22.0	8.6
18.5-20.0	14.5	8.9
20.0-25.0	34.1	6.5
>25.0	42.7	4.4
Standard of Living Index		
Low	15.5	12.1
Medium	25.7	7.0
High	38.9	5.0

Table 9: Percentage of Females Whose PPA Less Than Four Months (And Breastfeeding Duration Is More Than Six Months) and Median Duration of PPA According to Different Characteristics, **Karnataka**, NFHS-2

Characteristics	Percentage of females (whose PPA<4months, when BF>6)	Median duration of PPA (in months)
Total	19.0	8.2
Caste		
SC/ST	13.6	12.0
OBC	18.9	8.1
Others	22.8	6.8
Religion		
Hindu	17.5	8.3
Muslim	22.4	8.1
Place of Residence		
Rural	16.2	8.6
Urban	25.5	6.6
Education		
Illiterate	14.2	10.2
Literate (less than middle)	20.7	8.3
Middle to high school	19.7	6.8
Above high school	28.5	6.3
Body Mass Index		
<18.5	15.6	8.9
18.5-20.0	17.3	8.4
20.0-25.0	23.3	6.7
>25.0	29.6	6.3
Standard of Living Index		
Low	12.4	12.2
Medium	19.4	7.0
High	31.9	6.1

Table 10: Percentage of Females Whose PPA Less Than Four Months (And Breastfeeding Duration Is More Than Six Months) and Median Duration of PPA According to Different Characteristics, **Rajasthan**, NFHS-2

Characteristics	Percentage of females (whose PPA<4months, when BF>6)	Median duration of PPA (in months)
Total	33.8	6.9
Caste		
SC/ST	27.7	8.9
OBC	30.3	10.2
Others	40.0	6.2
Religion		
Hindu	33.8	6.9
Muslim	30.9	12.0
Place of Residence		
Rural	31.4	9.0
Urban	44.3	4.6
Education		
Illiterate	30.0	8.8
Literate (less than middle)	38.3	6.1
Middle to high school	48.9	4.1
Above high school	58.0	2.7
Body Mass Index		
<18.5	31.0	8.8
18.5-20.0	34.7	7.0
20.0-25.0	34.6	6.6
>25.0	55.2	2.6
Standard of Living Index		
Low	27.5	8.9
Medium	31.7	8.6
High	46.9	4.3

Table 11: Percentage of Females Whose PPA Less Than Four Months (And Breastfeeding Duration Is More Than Six Months) and Median Duration of PPA According to Different Characteristics, **Orissa**, NFHS-2

Characteristics	Percentage of females (whose PPA<4months, when BF>6)	Median duration of PPA (in months)
Total	23.2	8.9
Caste		
SC/ST	17.8	12.0
OBC	24.0	8.7
Others	31.6	6.5
Religion		
Hindu	23.4	8.8
Muslim	12.5	14.1
Place of Residence		
Rural	22.5	8.9
Urban	29.5	8.2
Education		
Illiterate	26.2	10.4
Literate (less than middle)	25.9	8.0
Middle to high school	30.6	6.6
above high school	40.7	4.8
Body Mass Index		
<18.5	21.7	10.4
18.5-20.0	24.2	8.9
20.0-25.0	25.6	6.9
>25.0	44.5	4.5
Standard of Living Index		
Low	19.1	12.0
Medium	25.5	8.1
High	39.4	5.0

Table 12: Percentage of Females Whose PPA Less Than Four Months (And Breastfeeding Duration Is More Than Six Months) and Median Duration of PPA According to Different Characteristics, **Uttar Pradesh**, NFHS-2

Characteristics	Percentage of females (whose PPA<4months, when BF>6)	Median duration of PPA (in months)
Total	32.1	8.8
Caste		
SC/ST	27.1	12.1
OBC	29.0	10.4
Others	35.5	8.0
Religion		
Hindu	35.6	8.5
Muslim	35.2	7.0
Place of Residence		
Rural	29.1	10.8
Urban	40.7	6.2
Education		
Illiterate	27.2	12.1
Literate (less than middle)	35.6	6.6
Middle to high school	40.5	6.7
Above high school	50.6	3.0
Body Mass Index		
<18.5	30.4	10.4
18.5-20.0	30.0	9.0
20.0-25.0	37.6	6.8
>25.0	54.5	2.8
Standard of Living Index		
Low	24.9	12.2
Medium	31.0	8.7
High	47.9	4.3

Table 13: Percentage of Females Whose PPA Less Than Four Months (And Breastfeeding Duration Is More Than Six Months) and Median Duration of PPA According to Different Characteristics, **Andhra Pradesh**, NFHS-2

Characteristics	Percentage of females (whose PPA<4months, when BF>6)	Median duration of PPA (in months)
Total	17.1	8.3
Caste		
SC/ST	12.7	10.2
OBC	16.2	8.5
Others	22.8	6.6
Religion		
Hindu	15.6	8.6
Muslim	35.4	4.9
Place of Residence		
Rural	15.7	8.6
Urban	21.2	6.7
Education		
Illiterate	11.0	10.1
Literate (less than middle)	23.3	8.1
Middle to high school	22.6	8.0
Above high school	26.5	4.8
Body Mass Index		
<18.5	14.0	10.2
18.5-20.0	15.4	8.3
20.0-25.0	24.1	6.3
>25.0	24.3	6.3
Standard of Living Index		
Low	11.1	10.7
Medium	18.5	8.1
High	27.8	4.8

Table 14: Percentage of Females Whose PPA Less Than Four Months (And Breastfeeding Duration Is More Than Six Months) and Median Duration of PPA According to Different Characteristics, **Madhya Pradesh, NFHS-2**

Characteristics	Percentage of females (whose PPA<4months, when BF>6)	Median duration of PPA (in months)
Total	26.7	10.7
Caste		
SC/ST	17.6	12.4
OBC	30.8	8.6
Others	40.0	6.4
Religion		
Hindu	30.8	8.4
Muslim	36.8	6.7
Place of Residence		
Rural	23.5	12.1
Urban	36.6	7.4
Education		
Illiterate	21.9	12.3
Literate (less than middle)	27.8	8.9
Middle to high school	38.0	6.2
above high school	48.2	4.2
Body Mass Index		
<18.5	23.4	12.1
18.5-20.0	27.1	10.6
20.0-25.0	29.1	10.2
>25.0	48.9	4.5
Standard of Living Index		
Low	20.6	12.4
Medium	24.8	12.0
High	44.0	4.9

Table 15: Percentage of Females Whose PPA Less Than Four Months (And Breastfeeding Duration Is More Than Six Months) and Median Duration of PPA According To Different Characteristics, **Bihar, NFHS-2**

Characteristics	Percentage of females (whose PPA<4months, when BF>6)	Median duration of PPA (in months)
Total	32.4	6.1
Caste		
SC/ST	24.7	12.3
OBC	28.7	8.0
Others	39.7	5.8
Religion		
Hindu	31.1	7.0
Muslim	43.1	4.4
Place of Residence		
Rural	26.6	10.1
Urban	41.5	4.8
Education		
Illiterate	24.0	10.7
Literate (less than middle)	34.9	8.2
Middle to high school	38.3	6.0
above high school	42.0	4.7
Body Mass Index		
<18.5	24.5	10.5
18.5-20.0	27.6	10.2
20.0-25.0	42.8	4.8
>25.0	38.4	6.1
Standard of Living Index		
Low	19.5	12.5
Medium	34.3	6.6
High	44.6	4.4

References

- [1] Akin J.S., Billsborrow R. E. Guilkey D. K. and Popkin B. M. (1986): "Breast-feeding Patterns and Determinants in the Near East: An Analysis for four countries" Population Studies. vol. 40 pp 247-262. <https://doi.org/10.1080/0032472031000142056>.
- [2] Bongaarts J. (1978): Framework for analysing the proximate determinants of fertility. Population Development Review vol. 4, pp.105-132. <https://doi.org/10.2307/1972149>.
- [3] Bongaarts J. and Delgado H. (1979): Effects of nutritional status on fertility in rural Guatemala. In: J.A.Menken and H. Lerdidon(eds),Patterns and Determinants of natural fertility, Liege, Ordina, pp. 107-033.
- [4] Bongaarts J. (1980): Does malnutrition affect fecundity? A summary of evidence. Science, vol.208 pp.564-569. <https://doi.org/10.1126/science.7367878>.
- [5] Chevez A. and Matinez C. (1973): Nutrition and development of infants from poor rural area -Maternal nutrition and its consequences on fertility. Nutr. Rep. Int., vol. 7, pp. 1-4
- [6] Chowdhury A.K.M.A. (1978): Effect of maternal nutritional status on fertility in rural Gwatemala. In: W.H. Mosley (ed.) Nutiation and human reproduction New York: pienum, pp. 401-409.
- [7] Delgado H., Lechtig A., Martorell R., Brineman E. and Klein R.E. (1978): Nutration, location and postpartum amenorrhoea. American Journal of Clinical Nutrition, vol. 31, pp.322-327.
- [8] Habicht,J.P. Davanzo, J.S., Butz, W.P. and Meyers, L. D., (1985). The contraceptive role of breastfeeding. Population Studies, vol. 39 pp.213-232. <https://doi.org/10.1080/0032472031000141456>.
- [9] Howie P.W. and Mc Neilly A.S.(1982) Contraceptive effect of breast-feeding patterns on Human Birth Intervals.Journa; of Reproduction and Fertility vol.65, pp.545
- [10] Huffman S.L., Chowdhury A.M.K., Chakraborty J. and Mosley W.H. (1978): Nutrition and post partum amenorrhoea in rural Bangladesh. Population Studies, vol. 32 pp. 251-260. <https://doi.org/10.1080/00324728.1978.10410713>.
- [11] Huffman S.L., Ford K., Allen H.A. and Streble, P. (1987) Nutrition and Fertility in Bengladesh: Breastfeeding and post partum amenorrhoea. Population Studies, vol. 41, pp.447-462. <https://doi.org/10.1080/0032472031000142996>.
- [12] ICMR (1995): Lactation-current concepts and concerns ICMR Bulletin, vol. 25 (8), pp.73-80
- [13] Nath D. C., Land K. C. and Singh K. K. (1994): The role of breast-feeding beyond postpartum amenorrhoea on the return of fertility in India: a life table and hazards model analysis Journal of Biosocial Science vol. 26, pp.191-206. <https://doi.org/10.1017/S0021932000021234>.
- [14] Nath D. C., Singh K.K., Land K. C.Talukdar (1993):Breastfeeding and PPA in traditional society: A Hazards modal analysis. Social Biology vol.40, pp.74-86.
- [15] National Family Health Survey 1998-1999, International Institute for Population Sciences, Mumbai, India.
- [16] Prema K., Naidu N., Neelakumari S. and Ramalakshmi, B.A. (1981): Nutrition-fertility interaction in lactating women of low-income groups. American Journal of Clinical Nutrition, vol. 45, pp.461-467. <https://doi.org/10.1079/bjn19810124>.
- [17] Santow,G. (1987):Reassessing the cotraceptive effect of breastfeeding.population studies vol.41,pp.147
- [18] Savina G. and Kennedy K. I. (1989): The effect of a breast-feeding education program on loctational amenrrhoea in the Philippines studies in Family planning vol 20 pp203-214.
- [19] Singh K. K., Suchindran C. M. and Singh R. S. (1994): Smoothed Breastfeeding duration and waiting time: Social Biology vol. 41(3,4), pp. 229-239.
- [20] Singh S. N. and Singh K. K. (1989): Life table analysis of censored data on post partum amenorrhoea period. Demography India, vol. 18(1-2), 27-38.
- [21] Singh S. N., Singh K. K, Singh K.(1990):Socio-economic development and transition in the duration of PPA. Survival function analysis of data, Janasamkhyia vol. (8) pp. 41-54.
- [22] Srinivasan K., Pathak K. B. and Pandey A. (1989): Determinants of breast-feeding and post partum amenorrhoea in Orissa. Journal of Biosocial Science vol. 21, pp. 365-371. <https://doi.org/10.1017/s002193200001806x>.
- [23] Yadav S., Mishra C.P. and Srivastava P. (2005): Impact of Breast-feeding vis-à-vis Nutrition on Duration of Post Partum Amenorrhoea, Demography India, vol. 34 (2), pp. 233-239.